

Thesis on Peritonitis
for a
Degree of M. D.
by
James H. Gilliam
of Va.

Deposited March 16th 1874
W. S. H
Dean

of us
he a
comp
he c
est
treat
later
re of
land
tack
ly ar
ing
part
Wh
is
to the
myon

In obedience to the requisitions for graduation I have selected a view of Peritoneal inflammation for discussion:

Though from the limited extent of observation and experience of the noviciate in medicine no novel idea can be reasonably expected, yet our attempt will be to offer such an analysis of the subject as will manifest in some degree an acquaintance with the general principles of science.

This result being the proper purpose of inaugural disquisitions, I confidently appeal for approbation to that liberality of sentiment which distinguishes the illustrious Professors,

[Faint, illegible handwriting, likely bleed-through from the reverse side of the page.]

His
re m
appe
al m
tious
the
who ar
Ch
melt
nd e
whole
t ab
the o
stern
which
of he
bela
he fac
the p
refr

of this celebrated University-

Of the inflammations of the abdominal viscera Peritonitis is comparatively a new disease, hence the contrariety of opinion with respect to its most successful mode of treatment.

All agree that it is most fatal in its tendency, and of all others one of the most insidious in its approach and progress.

When this affection attacks the male sex it almost invariably arises in consequence of inflammation being translated from some other part, as, the stomach, liver &c.

When the other sex is the subject of it, it is most generally consecutive to the parturient condition, and thereby conjoined with the puerperal fever.

[Faint, illegible handwriting on lined paper, likely bleed-through from the reverse side.]

...nd a
...py
...nd n
...turt
...hoal
...milit
...ome
...nour
...ous
...m
...be
...pro
...ale
...fig
...ab
...le
...th
...as
...not

Its most common and efficient causes, are suppression of perspiration, the application of cold to the surface, mechanical injuries, as contusion, severe exercise, tedious and difficult labours, &c.

Like the other Phlegmasia it is divided into an acute and chronic stage.

In the acute stage the inflammation sometimes attacks only a small part, and is subsequently imparted to the whole of it; but most commonly it attacks the whole at once.

The onset of the disease is usually attended with chills and shiverings, which are soon followed by a diffusion of heat over the body, and pain in the abdomen more or less so extensive as to be felt over the whole of that part. The pain is greatly augmented by pressure. The pulse is frequent, small

for per
ances
the bu
the pa
d is m
manuf
piece
same
which
Dark
spinn
cold c
tis ar
is haq
fiend
the see
tong th
of inf

and contracted, the skin becomes hot & dry, the tongue is parched and encrusted and now the patient complains of much thirst. Even in the early stage the tongue, throat and fauces assume an appearance similar to that in typhus fever, tho' sometimes they are found perfectly moist.

In the course of ten or twelve hours, the tenderness of the abdomen is so much increased, that the weight of the bed clothes becomes insupportable. The pulse becomes greatly contracted and beats from one hundred to one hundred and fifty in a minute, and the swelling of the abdomen becomes apparent. The patient lies constantly on his back with his knees elevated or drawn up so as to relax the abdominal muscles, and this being a peculiar symptom, should

infr
lone
Dagn
inter
sym
fatu
he fo
mole
from
and
from
Dagn
omen
life
cole
reap
and
pair
on

be particularly noticed.

As the disease advances, all these symptoms, particularly the tymefaction, increase. But occasionally the pain suddenly subsides, which, when it is not the effect of our remedies, is an inauspicious omen, being always the precursor of death. There is about the same time a great swelling of the joints, which is very much accelerated.

Dark matter is vomited, Singultus, or spasmodic affections of the stomach and cold clammy sweats occur. The extremities are cold and withered, the countenance is haggard and collapsed and at last difficult and laborious respiration closes the scene.

We find by postmortem examinations the peritoneum exhibiting the stages of inflammation, and the bowels not

For
last
year

area
in

of the
with

after

and

for

also

robust

new

by

by

our

my

gate

ing

infrequently in such a state as to amount almost to gangrene.

Diagn. The Dentition very much resembles miteris and Cholera in its more general symptoms, yet it assumes some peculiar features in its character.

It differs from the former by the pain being permanent, by producing no desire to go to stool, and from both by the position of the patient and the slightest alleviation not arising from the most copious alvine evacuation.

Progn. The following are favourable omens, the pulse becoming fuller and less frequent, the skin moister and cooler, the perspiration less laborious, the reappearance of the lochial discharge and the gradual diminution of the pain and tension of the abdomen. On the other hand the sudden ceasing



tion of pain with oppression and tumefaction are fatal symptoms.

Treatment—

In the removal of this disease nearly the same mode of cure is indicated as in that of inflammation of the Menstrua. We attack the acute form with venerection to ^{the} greatest practicable extent as the most powerful evacuant, and extend it, regardless of the pulse, as far as the strength of the patient will allow. If the patient be adult of a robust constitution, we often find it necessary to take from twenty to forty ounces at one bleeding, and this quantity should even be repeated on the course of ten or twelve hours, if the symptoms are not materially mitigated. If after the lancet has been urged to this extent, some of the most



prominent symptoms still remain, as
 pain continue we should resort to
 cups or leeches, and remove as much
 blood as can be effected by these means.
 Emollient fomentations or warm poultices
 applied to the abdomen sometimes
 produce considerable relief. The best
 application is a bread and milk poultice
 in a sack moderately filled with warm
 milk so as to accommodate itself to the
 form of the abdomen. By a Writer of no
 inconsiderable authority it is contended that
 poultices are as well adapted to acute sea-
 red inflammations as to those situated
 on the surface. On this as it may, they
 often abridge pain and make a favora-
 ble impression. When blood letting can
 be no longer used diaphoretics consti-
 tute an important part of the curative
 plan. By their centrifugal tendency

at the
low
little
light
bride
my
men
Altho
pious
yet
in a
milit
the la
magn
large
apost
of the
obli
in y

they lessen or entirely remove deep seated congestions, and produce on the surface a most salutary effect. With a particular view to induce perspiration we resort to external applications, and of these the vapour bath merits the most confidence. To increase and assist its impression we should give the Dover's powder, or some other preparation, of which opium constitutes the principal part. The vapour bath has been beneficially used, and when considered necessary should not be neglected.

Whatever opposition may have been advanced against blisters we repose considerable confidence in their utility. When timely applied in this case as in the Phlegmona generally, they not infrequently extinguish the remaining pain and avert the progress of the disease. The proper time of their application we consider to be when

of 200
Tons
The do
house
Winds
Was r
some
calcu
In a
grea
on the
prod
malt
depre
is to
dura
draw
tract
arous
by 10

after general and local depletion has
been used to a considerable extent, there
will remain a permanent of pain and
slight degree of tenderness to the touch.
Under these circumstances blisters of a
sufficient size to cover the whole abdo-
men produce the most beneficial effects.
Altho we have asserted that evacuations
from the bowels do not abate the pain,
yet we deem it necessary to keep them
in a soluble state. For this purpose
mild laxatives or enemata will answer;
the latter should be composed of mild
ingredients and administered in such
large quantities as to act as emollient
applications to the parts.

When in despite
of these remedies the disease continues,
obstinate and appears about to terminate
in gangrene our best recourse is to Ept.

for a
the s
the the
male
to est
ten a
milla
chara
as he
stann
being
men
ng be
a 2.5
This
apian
my m
the s
the i

of impeture which has often asserted this tendency, when all other remedies have failed. The dose is a table spoonful every two or three hours. The treatment detailed above is much adapted to an ordinary attack of this disease; but there are occasionally some variations in the symptoms well calculated to mislead the practitioner.

In its early stage there is sometimes so great a prostration of strength attended with a pulse so extremely feeble as to produce an apprehension of immediate death. This we observe a case of extreme depression in which the energy of the body is so overwhelmed by the force of the disease, that if blood should be copiously drawn the system would be unable to react. Under this supposition we strive to arouse the depleted energies of the system by the warm bath, naphoretics and small



bleedings, watching with great caution
the state of the inside while the blood flows
in this way by cautiously coagulating away
small portions of blood we make a flaw
to establish a free and general circula-
tion, and thus develop an acute state of
inflammation to be ^{managed} as previously directed.

There is also a case of an opposite
character in which the disease advances
so treacherously as to excite no particular
alarm in the patient or his friends, there
being only a slight tenderness of the abdo-
men, very little fever and a pulse differ-
ing from its healthy state only in being
a little quicker and somewhat chorded.

This delusive appearance is soon made
apparent by the succession of a train of
symptoms of the most alarming kind.
The system becomes suddenly depressed,
the pulse sinks, the surface is covered with

4

B

my
from
to be
to the
from
of ge
mple
boma
boma
of ge
are
to n
aner
late
lion

a cold clammy sweat, and the patient with-
out the aid of the most active remedies,
directed by the mercurial judgment merit-
ly sinks.


The chronic form of this disease
also approaches very insidiously. It commences
with a slight soreness of the abdomen
and more or less accelerated, the tongue
in the morning is furred and there is some
complaint of thirst. The face is pale, the
features are a little shrunk with a slight
diminution of temperature in the extremi-
ties. These symptoms continue sometimes
without creating any alarm, until by some
exciting cause the disease assumes an
acute form. And when this occurs
it becomes the most terrible of all diseases.
Here we are debarred from the use of the
same as the same extent as in diseases




originally acute, and the ~~refers~~ are so strongly perverted by wrong action, as to be extremely difficult of restoration to their former state.

The same mode of cure is pursued here except the extent of general bleeding, therefore that by cups or leeches is more particularly demanded, to be followed by fomentations, blisters and in emergencies to the Spirits of Turpentine.

This form of the disease as soon as it is discovered, should be managed by frequent small bleedings, general and local, observing a complete state of rest and a most rigid restriction to the antiphlogistic regimen.





101

Page 101
H. 1. 11
C. 1. 11

for
supplied
the
may
making

at the
all
of
the
be a
at as

under
order
Ent
the in
found